不同粒径聚苯乙烯微塑料和苯并(a)芘联合暴露影响炎症因子的表达

徐雅雯, 朱军, 胡堇叶, 张展, 李磊, 吴倩*

南京医科大学公共卫生学院卫生检验学系, 南京 211166

* 联系人, E-mail: wuqian@njmu.edu.cn

2020-06-05 收稿, 2020-08-14 修回, 2020-08-14 接受, 2020-08-17 网络版发表

江苏省自然科学基金(BK20161571)、江苏省高校自然科学研究重大项目(16KJA330002)、江苏高校品牌专业建设工程资助项目(PPY2015A067)和江苏高校优势学科建设工程项目资助

摘要 研究发现, 微塑料可吸附并富集持久性有机污染物, 吸附污染物后的微塑料生物毒性增强. 本文旨在探究不同粒径聚苯乙烯微塑料(polystyrene microplastics, PS-MPs)联合苯并(a)芘(benzo(a)pyrene, BaP)暴露, 对成年斑马鱼鳃和支气管相关细胞炎症细胞因子表达的影响. 以两种粒径的PS-MPs(0.5和5 μm)分别与BaP联合暴露幼鱼7 d、成鱼14 d和细胞48 h. 使用Real-time PCR测定不同暴露组炎症因子基因相对表达情况, 体视镜观察微塑料颗粒进入幼鱼体内的情况, 共聚焦显微镜观察微塑料颗粒进入细胞的情况. 结果发现, 粒径为0.5和5 μm 的PS-MPs均能进入幼鱼体内; 0.5 μm PS-MPs单独暴露时, *TNF- α* 、*IL-6*和*IL-8*基因相对表达显著高于对照组; 5 μm PS-MPs单独暴露时, *IFN- γ* 基因相对表达显著高于对照组. BaP和0.5 μm PS-MPs联合暴露时, 成鱼鳃组织*IFN- γ* 、*IL-6*和*IL-8*基因的相对表达高于BaP单独暴露. 细胞实验结果表明, 仅0.5 μm 的PS-MPs能进入两种细胞胞质中. 在人支气管上皮样细胞HBE中0.5 μm PS-MPs组炎症因子*TNF- α* 、*IL-6*和*IL-8*基因相对表达均高于对照组. 在人支气管平滑肌细胞HBSMC中0.5 μm PS-MPs组炎症细胞因子*TNF- α* 、*IL-8*和*IL-1 β* 的相对表达显著高于对照组. HBE细胞中BaP分别与0.5 μm /5 μm 微塑料联合暴露时, 均未发现明显的联合作用. HBSMC细胞中BaP和0.5 μm PS-MPs联合暴露时, 对*IFN- γ* 和*IL-6*炎症因子的表达有显著的联合作用; 而BaP和5 μm PS-MPs联合暴露时, 有显著减弱BaP促炎作用的趋势, 特别是对*TNF- α* 和*IL-8*炎症因子表达的影响. 由此推断, 粒径可能是影响微塑料生物摄取和累积的主要因素之一, 0.5 μm 微塑料因其易被细胞摄取, 故联合有机污染物暴露可加重炎症反应. 另外, 因5 μm 微塑料未能进入细胞, 其对炎症反应的影响仍需进一步探究.

关键词 聚苯乙烯微塑料, 苯并(a)芘, 炎症, 斑马鱼, 细胞

微塑料指直径<5 mm的塑料颗粒^[1], 比表面积大和疏水性的特点使其更容易吸附污染物, 能够作为许多有害化学物质的载体, 产生复合毒性效应. 有报道, 微塑料可以富集海洋环境中的有机污染物^[2], 增强其在海洋生物体内的生物效应, 包括免疫反应、氧化应激反应等^[3]. 近年来, 微塑料的研究已经从海洋转向陆地和

大气^[4,5], 有研究人员在肺癌患者肺部中发现了吸入纤维素和塑料纤维^[6], 说明微塑料可以通过大气“旅行”进入呼吸道^[7]. 然而, 目前大气微塑料吸入及联合有机污染物暴露对人体的健康风险研究较少, 且致病机制尚不明确, 因此开展相关研究至关重要.

随着经济的快速发展, 燃煤取暖和汽车尾气排放

引用格式: 徐雅雯, 朱军, 胡堇叶, 等. 不同粒径聚苯乙烯微塑料和苯并(a)芘联合暴露影响炎症因子的表达. 科学通报, 2020, 65: 4281–4290

Xu Y W, Zhu J, Hu J Y, et al. Co-exposure to different sized polystyrene microplastics and benzo[a]pyrene affected inflammation in zebrafish and bronchial-associated cells (in Chinese). Chin Sci Bull, 2020, 65: 4281–4290, doi: [10.1360/TB-2020-0668](https://doi.org/10.1360/TB-2020-0668)

等因素使得多环芳烃(polycyclic aromatic hydrocarbon, PAHs)成为城市空气中不容忽视的重点污染物之一。流行病学研究显示, PAHs暴露与儿童呼吸系统疾病如支气管炎发生发展相关^[8], 并影响成人肺功能^[9]。多环芳烃由细胞色素P450^[10]代谢形成二醇-环氧化物, 与细胞大分子共价结合, 诱导PAH-DNA加合物的形成, 其具有细胞毒性、基因毒性, 可引起肺部炎症、甚至致癌^[11]。微塑料能够吸附亲脂性PAHs^[12], 但二者联合暴露对呼吸系统造成的潜在健康风险机理尚不清晰。本研究拟采用斑马鱼模型和支气管相关细胞模型对微塑料颗粒联合苯并(a)芘暴露是否加重气道炎症反应进行初步探究。

1 材料和方法

(i) 实验材料。成年野生型斑马鱼购自南京尧顺禹生物科技有限公司, 鱼龄为6个月。斑马鱼饲料购自山东升索渔用饲料研究中心。人支气管上皮样细胞(HBE)和人支气管平滑肌细胞(HBSMC)由南京医科大学公共卫生学院储海燕副教授惠赠。胎牛血清和高糖培养基(DMEM)购于美国Gibco公司。活细胞染色液(Hoechst33342)和细胞膜红色荧光探针(DiI)购自上海碧云天生物技术有限公司。苯并(a)芘(Benzo[a]pyrene, $\geq 96\%$, HPLC)购自美国Sigma公司(1 mg/mL储备液)。MS-222试剂购自南京尧顺禹生物技术股份有限公司。聚苯乙烯绿色荧光微球购自天津倍思乐色谱技术开发中心, 浓度10 mg/mL(工作浓度1 mg/L), 直径分别为0.5和5 μm , 最大发射波长为518 nm, 最大激发波长为488 nm。DNA引物购自南京金斯瑞生物科技有限公司。

(ii) 斑马鱼实验。将成鱼随机分为6组: 对照组、1 mg/L 0.5 μm 微塑料组^[13-15]、1 mg/L 5 μm 微塑料组、10 $\mu\text{g/L}$ BaP组、BaP和0.5 μm 微塑料联合暴露组、BaP和5 μm 微塑料联合暴露组, 每组9条(雌鱼6条, 雄鱼3条)。在28°C环境, 14:10明暗光周期条件下, 人工饲料辅以丰年虫, 每日2次, 未通氧。为防止塑料颗粒沉积, 培养液隔天更换一次, 共培养14 d。剖杀取两侧鳃组织, 液氮保存。为观察微塑料在斑马鱼中的分布情况, 将成年雌雄斑马鱼按1:1配对产卵, 暴露于加入上述浓度微塑料的培养液中, 培育7 d, 每天更换培养液, 第7天时使用体视荧光显微镜SMZ18(Nikon, 日本)观察微塑料进入幼鱼体内情况。

(iii) 细胞实验。HBE和HBSMC用含10%胎牛血清

DMEM高糖完全培养基置于37°C, 5%CO₂培养箱中培养。细胞分为6组: 对照组、50 $\mu\text{g/mL}$ 0.5 μm 微塑料组、50 $\mu\text{g/mL}$ 5 μm 微塑料组、1 nmol/L BaP组、BaP和0.5 μm 微塑料联合暴露组、BaP和5 μm 微塑料联合暴露组。37°C培养48 h, 依据说明书方法分别将核与胞质染色后, 使用激光共聚焦显微镜700B(Zeiss, Germany)拍照, 同时收集细胞, 液氮保存。

(iv) 炎症因子检测。检测的炎症因子包括TNF- α 、IFN- γ 、IL-6、IL-8和IL-1 β 。冻存的斑马鱼鳃组织和细胞, TRIzol(Invitrogen, 美国)法提取总RNA, 核酸蛋白分析仪(Thermo NanoDrop 2000, 美国)检测浓度。反转录使用试剂盒PrimeScriptTM RT reagent Kit(TaKaRa Bio Inc, 日本), 具体操作根据说明书。PCR体系包含12.5 μL 2 $\times$ SYBR Green II Buffer(TaKaRa Bio Inc, 日本), 1 μL 正向引物(10 $\mu\text{mol/L}$), 1 μL 反向引物(10 $\mu\text{mol/L}$)和2 μL cDNA。PCR反应参数: 95°C变性10 min, 然后在95°C变性5 s, 60°C退火延伸45 s, 共40个循环。每个循环在60°C获得荧光数据。采用LightCycler 96(Roche, 瑞士)进行实时定量荧光PCR检测分析。引物设计使用Primer Premier 5.0, 如表1所示。GAPDH作为细胞基因表达内参, β -catin作为斑马鱼基因表达内参, 设置3个平行。采用 $2^{-\Delta\Delta C_t}$ 法处理数据, 相对表达计算公式:

$$\Delta C_t = \text{Mean}C_{t(\text{测量基因})} - \text{Mean}C_{t(\text{GAPDH基因})};$$

$$\Delta\Delta C_t = \Delta C_t(\text{实验组}) - \Delta C_t(\text{对照组}) \quad (1)$$

(v) 数据统计分析。采用*t*检验比较各暴露组与对照组之间差异性; 单因素方差分析比较各暴露组间差异性, $P < 0.05$ 具有统计学意义。采用GraphPad Prism 6.0软件分析和作图。

2 结果

2.1 斑马鱼幼鱼体内微塑料分布情况

如图1所示, 粒径为0.5和5 μm 的微塑料颗粒均能进入幼鱼体内, 实验组的幼鱼胃肠道内可观察到绿色荧光颗粒; 对照组的幼鱼体内无微塑料颗粒。

2.2 微塑料暴露影响成年斑马鱼鳃中炎症细胞因子的相对表达

如图2所示, 0.5 μm 微塑料组的TNF- α 、IL-6和IL-8基因相对表达显著高于对照组; 5 μm 微塑料组的IFN- γ 基因相对表达显著高于对照组。

表 1 PCR引物序列

Table 1 PCR primer sequences

基因名称	上游引物序列	下游引物序列
<i>Danio rerio</i> β -actin	5'-GTGACCGTCCAGAAATGCTT-3'	5'-CACCAGGACCACTCCCTAAA-3'
<i>TNF-α</i>	5'-CAAAACATTCTTCGGGGTGT-3'	5'-ACTCTCACTGCATCGGCTTT-3'
<i>IFN-γ</i>	5'-CTTCAGACAACCAGCGCATA-3'	5'-CAGCATTCCTTCAAGCAACA-3'
<i>IL-6</i>	5'-TCTGCCACACCTGAAGACAG-3'	5'-GAAGCCGAGTTAGGACGATG-3'
<i>IL-8</i>	5'-CTACCGAGACGTGGGTGATT-3'	5'-CGTGTGTGTGAGGAGGGAAGT-3'
<i>Homo sapiens</i> GAPDH	5'-GAAGGTGAAGGTCCGAGTC-3'	5'-GAAGATGGTGATGGGATTTC-3'
<i>TNF-α</i>	5'-CCCTCACACTCACAAACCAC-3'	5'-ACAAGGTACAACCCATCGGC-3'
<i>IFN-γ</i>	5'-GCGCCAAGCATTCAATGAGC-3'	5'-ATCTCTTCCCCACCCGAAT-3'
<i>IL-6</i>	5'-CCTTCGGTCCAGTTGCCTTCTCCCT-3'	5'-GGGCTGAGATGCCGTCGAGGATGTA-3'
<i>IL-8</i>	5'-CCACCGGAAGGAACCATCTC-3'	5'-TTCCTTGGGGTCCAGACAGA-3'
<i>IL-1β</i>	5'-ACCACTACAGCAAGGGCTTCAGGCA-3'	5'-TCCTGTCCCTGGAGGTGGAGAGCTT-3'

图 1 体视荧光显微镜下斑马鱼幼鱼中微塑料分布. 绿色荧光颗粒为聚苯乙烯塑料微粒. (a)~(c) 正常光照片; (d)~(f) 最大发射波长518 nm时的荧光照片. (a), (d) 对照组; (b), (e) 0.5 μ m微塑料组; (c), (f) 5 μ m微塑料组

Figure 1 PS-MPs distribution in zebrafish by stereoscopic fluorescence microscope. The green fluorescent particles are PS-MP particles. (a)~(c) Normal photos; (d)~(f) fluorescence photos with the maximum emission wavelength of 518 nm. (a), (d) Control group; (b), (e) 0.5 μ m MP group; (c), (f) 5 μ m MP group

2.3 BaP和微塑料联合暴露影响成年斑马鱼鳃中炎症细胞因子的相对表达

如图3所示, BaP与0.5 μ m微塑料联合暴露组 $IFN-\gamma$ 、 $IL-6$ 和 $IL-8$ 基因的表达显著高于BaP单独暴露组; BaP与5 μ m微塑料联合暴露组仅 $IL-6$ 基因的表达显著高于BaP单独暴露组, 提示BaP和0.5 μ m微塑料联合暴

露时产生明显的联合作用.

2.4 微塑料的粒径影响HBE细胞和HBSMC细胞摄取

如图4所示, 粒径为0.5 μ m的微塑料颗粒被摄入并分布在两种细胞胞质中; 粒径为5 μ m的微塑料颗粒未进

图 2 不同粒径微塑料暴露对斑马鱼鳃炎症性细胞因子基因相对表达的影响. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

Figure 2 Effects of PS-MPs exposure with different particle sizes on relative expression of inflammatory factors in zebrafish gill. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

图 3 BaP与微塑料联合暴露对斑马鱼鳃炎症性细胞因子基因相对表达的影响. * $P < 0.05$, **** $P < 0.0001$

Figure 3 Effects of co-exposure with PS-MPs and BaP on relative expression of inflammatory factors in zebrafish gills. * $P < 0.05$, **** $P < 0.0001$

图4 激光共聚焦显示微塑料在HBE细胞和HBSMC细胞内的分布情况。镜下胞质呈红色、核呈蓝色、聚苯乙烯微塑料为圆形绿色荧光颗粒。
(a) HBE细胞0.5 μm微塑料组; (b) HBSMC细胞0.5 μm微塑料组; (c) HBE细胞5 μm微塑料组; (d) HBSMC细胞5 μm微塑料组

Figure 4 The distribution of PS-MPs in HBE and HBSMC cells by laser confocal microscopy. The cytoplasm is red. The nucleus is blue. PS-MP are green fluorescent particles under laser confocal microscopy. (a) 0.5 μm MP in HBE; (b) 0.5 μm MP in HBSMC; (c) 5 μm MP in HBE; (d) 5 μm MP in HBSMC

入细胞内,说明粒径的尺寸影响细胞摄取微塑料颗粒。

2.5 微塑料暴露影响HBE细胞和HBSMC细胞中炎性细胞因子的相对表达

如图5所示,在HBE细胞中,0.5 μm微塑料组炎症因子*TNF-α*、*IL-6*和*IL-8*基因相对表达均高于对照组。如图6所示,在HBSMC细胞中,0.5 μm微塑料组炎症细胞因子*TNF-α*、*IL-8*和*IL-1β*的相对表达显著高于对照组。

2.6 BaP和微塑料联合暴露影响HBE细胞和HBSMC细胞中炎性细胞因子的相对表达

如图7所示,在HBE细胞中BaP分别和0.5 μm/5 μm

微塑料联合暴露组,均未发现明显的联合作用,不论是上调表达的*TNF-α*和*IL-8*炎症因子,还是下调表达的*IFN-γ*、*IL-6*和*IL-1β*炎症因子。如图8所示,在HBSMC细胞中,BaP和0.5 μm微塑料联合暴露组对*IFN-γ*和*IL-6*炎症因子的表达有显著的联合作用。而BaP和5 μm微塑料联合暴露组有显著减弱BaP促炎作用的趋势,特别是对*TNF-α*和*IL-8*炎症因子表达的影响。

3 讨论

在斑马鱼幼鱼实验中我们观察到粒径分别为0.5和5 μm的聚苯乙烯微塑料在鱼体内的分布和积累情况与相关文献结果基本一致。例如,Lei等人^[16]研究使用的

图5 不同粒径微塑料暴露对HBE细胞中炎性细胞因子基因相对表达的影响。* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$

Figure 5 Effects of PS-MPs exposure with different particle sizes on relative expression of inflammatory factors in HBE cells. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$

图 6 不同粒径微塑料暴露对HBSMC细胞中炎症细胞因子基因相对表达的影响. ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$
Figure 6 Effects of PS-MPs exposure with different particle sizes on relative expression of inflammatory factors in HBSMC cells. ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$

图 7 BaP与微塑料联合暴露对HBE细胞中炎症细胞因子基因相对表达的影响. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$
Figure 7 Effects of co-exposure with PS-MPs and BaP on relative expression of inflammatory factors in HBE cells. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

0.1、1和5.0 μm微塑料颗粒均能进入斑马鱼体内。Qiao等人^[17]的研究显示，5 μm的微塑料能够进入斑马鱼肠道。Batel等人^[6]的实验结果表明，1~5 μm粒径微塑料颗

粒在进入成年斑马鱼鳃中6和24 h后，大多数微塑料颗粒附着在鳃黏液层的纤维上，并不断被排出。

斑马鱼作为现代病理学、免疫学等研究中常用的

图 8 BaP与微塑料联合暴露对HBSMC细胞中炎症细胞因子基因相对表达的影响。* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

Figure 8 Effects of co-exposure with PS-MPs and BaP on relative expression of inflammatory factors in HBSMC cells. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

模式生物^[18]，与人类基因高度相似并具有高度保守的疾病信号转导通路^[19]。斑马鱼的先天和适应性免疫系统，包括巨噬细胞、粒细胞、Th1、Th2标记的T细胞和B细胞^[20]，肺部疾病相关的炎症细胞也与人类高度相似^[21,22]。斑马鱼鳃上皮的免疫细胞和平滑肌细胞分布在鳃板上，并有黏液覆盖^[23]，结构与哺乳动物的呼吸道相似，具有相同的气体交换功能，是研究肺部炎症的良好模型。

在斑马鱼实验中我们发现，0.5 μm微塑料组的*TNF-α*、*IL-6*和*IL-8*基因相对表达显著高于对照组；BaP与0.5 μm微塑料联合暴露组*IFN-γ*、*IL-6*和*IL-8*基因的表达显著高于BaP单独暴露组，产生了明显的联合作用。细胞实验中，在HBSMC细胞中，0.5 μm微塑料组炎症细胞因子*TNF-α*、*IL-8*、和*IL-1β*的相对表达显著高于对照组；BaP和0.5 μm微塑料联合暴露组对*IFN-γ*和*IL-6*炎症因子的表达有显著的联合作用，而BaP和5 μm微塑料联合暴露组有显著减弱对*TNF-α*和*IL-8*炎症因子表达的影响。我们认为，粒径可能是影响微塑料通过不同机制引发细胞炎症反应的因素之一：5 μm的聚苯乙烯微塑料颗粒无法进入细胞质中，仅在胞外造成机械损伤，引起细胞炎症反应；粒径为0.5 μm的微塑料被摄入胞内同时带入吸附于表面的BaP^[11]，破坏细胞生理

功能引发炎症并加重BaP对细胞的毒作用，如促进炎症细胞因子的表达升高。但Liu等人^[24]的研究结果显示，纳米聚苯乙烯微塑料颗粒尺寸的减小对吸附多环芳烃没有明显影响。本研究选择了微米级的微塑料，其中小粒径的为0.5 μm，对BaP的富集作用是否较粒径为5 μm的微塑料颗粒更强，有待进一步研究。

本研究主要检测了炎症细胞因子*TNF-α*、*IFN-γ*、*IL-1β*、*IL-6*和*IL-8*的相对表达情况。在斑马鱼和细胞模型中，BaP和PS-MPs联合暴露时炎症因子*IL-6*、*IL-8*表达的变化相对稳定。在炎症过程中，肿瘤坏死因子(tumor necrosis factor, *TNF*)、白细胞介素(interleukin, *IL*)和干扰素家族(interferon, *IFN*)等起主要作用。斑马鱼炎症因子*IL-8*与人类的具有高度同源性^[25,26]。*IL-8*由巨噬细胞和上皮细胞等分泌，与白细胞上的趋化因子受体结合并激活这些受体，诱导特异性细胞内信号传导级联，快速募集中性粒细胞^[27,28]，参与机体局部急性炎症初期过程^[29]，是各种炎症细胞所表达和释放的第一个信号^[30,31]。*IL-6*主要由单核巨噬细胞、Th2细胞、血管内皮细胞产生，诱导B细胞分化和产生抗体，诱导T细胞活化增殖、分化，参与机体免疫应答，是炎症反应的促发剂^[32]。我们推测，微塑料颗粒吸附BaP后进入呼吸道，通过支气管上皮细胞附着、摄入，进入肺泡并聚集

在终末细支气管中,引起肺巨噬细胞M1极化,而IL-6可通过对巨噬细胞的抑制作用进一步抑制Th1、Treg细胞的分化,促进Th2、Th17细胞的分化^[33],加重气道炎症反应。

另外,微塑料很大一部分来自微纤维。在生活中,以化纤为原料的物品非常普遍,如衣物、地毯等。衣物摩擦或在地毯上行走,会释放出微纤维进而形成微塑料。呼吸途径摄入微塑料可能更为直接,摄入量也非常大,可能与饮食摄入量相当,不容忽视。Dris等人^[34]报道,起居室或办公室内空气纤维类微塑料沉降通量可达 $1.60 \times 10^3 \sim 1.10 \times 10^4$ 个/(cm³ d)。本研究中仅采用聚苯乙烯微塑料及海洋环境中微塑料毒性作用浓度来分析

其生物效应具有一定局限性,与实际大气环境中微塑料种类和浓度可能不同,未来应聚焦研究大气环境不同种类、浓度和粒径的微塑料暴露的生物积累效应及联合持久性有机污染物后对人体的毒性效应。

4 结论

微塑料的粒径可能是影响生物摄取和累积的主要因素之一,联合有机污染物暴露可加重炎症反应。微塑料作为新兴的环境污染物之一,其自身和吸附其他环境污染物对人体健康产生的潜在危害值得关注。研究微塑料和大气污染物相互作用效应对探索其相互作用的机制和揭示微塑料危害人体呼吸系统有重要意义。

参考文献

- 1 Andrady A L. Microplastics in the marine environment. *Mar Pollut Bull*, 2011, 62: 1596–1605
- 2 Pauly J L, Stegmeier S J, Allaart H A, et al. Inhaled cellulosic and plastic fibers found in human lung tissue. *Cancer Epidemiol Biomarkers Prev*, 1998, 7: 419–428
- 3 Kim Y, Jeong J, Lee S, et al. Identification of adverse outcome pathway related to high-density polyethylene microplastics exposure: *Caenorhabditis elegans* transcription factor RNAi screening and zebrafish study. *J Hazard Mater*, 2020, 388: 121725
- 4 Liu M, Lu S, Song Y, et al. Microplastic and mesoplastic pollution in farmland soils in suburbs of Shanghai, China. *Environ Pollut*, 2018, 242: 855–862
- 5 Allen S, Allen D, Moss K, et al. Examination of the ocean as a source for atmospheric microplastics. *PLoS One*, 2020, 15: e0232746
- 6 Batel A, Borchert F, Reinwald H, et al. Microplastic accumulation patterns and transfer of benzo[a]pyrene to adult zebrafish (*Danio rerio*) gills and zebrafish embryos. *Environ Pollut*, 2018, 235: 918–930
- 7 Wright S L, Kelly F J. Plastic and human health: A micro issue? *Environ Sci Technol*, 2017, 51: 6634–6647
- 8 Hertz-Picciotto I, Baker R J, Yap P S, et al. Early childhood lower respiratory illness and air pollution. *Environ Health Perspect*, 2007, 115: 1510–1518
- 9 Cakmak S, Hebborn C, Cakmak J D, et al. The influence of polycyclic aromatic hydrocarbons on lung function in a representative sample of the Canadian population. *Environ Pollut*, 2017, 228: 1–7
- 10 Shimada T, Fujii-Kuriyama Y. Metabolic activation of polycyclic aromatic hydrocarbons to carcinogens by cytochromes P450 1A1 and 1B1. *Cancer Sci*, 2004, 95: 1–6
- 11 Pratt M M, John K, MacLean A B, et al. Polycyclic aromatic hydrocarbon (PAH) exposure and DNA adduct semi-quantitation in archived human tissues. *Int J Environ Res Public Health*, 2011, 8: 2675–2691
- 12 Batel A, Linti F, Scherer M, et al. Transfer of benzo[a]pyrene from microplastics to *Artemia nauplii* and further to zebrafish via a trophic food web experiment: CYP1A induction and visual tracking of persistent organic pollutants. *Environ Toxicol Chem*, 2016, 35: 1656–1666
- 13 Wegner A, Besseling E, Foekema E M, et al. Effects of nanopolystyrene on the feeding behavior of the blue mussel (*Mytilus edulis* L.). *Environ Toxicol Chem*, 2012, 31: 2490–2497
- 14 Lu Y, Zhang Y, Deng Y, et al. Uptake and accumulation of polystyrene microplastics in zebrafish (*Danio rerio*) and toxic effects in liver. *Environ Sci Technol*, 2016, 50: 4054–4060
- 15 Liu Z, Cai M, Yu P, et al. Age-dependent survival, stress defense, and AMPK in *Daphnia pulex* after short-term exposure to a polystyrene nanoplastic. *Aquat Toxicol*, 2018, 204: 1–8
- 16 Lei L, Wu S, Lu S, et al. Microplastic particles cause intestinal damage and other adverse effects in zebrafish *Danio rerio* and nematode *Caenorhabditis elegans*. *Sci Total Environ*, 2018, 619–620: 1–8
- 17 Qiao R, Sheng C, Lu Y, et al. Microplastics induce intestinal inflammation, oxidative stress, and disorders of metabolome and microbiome in zebrafish. *Sci Total Environ*, 2019, 662: 246–253
- 18 Chen Z L. Application of zebrafish in drug screens for active ingredients (in Chinese). *China J Chin Mater Med*, 2015, 40: 1235–1239 [陈志亮. 斑马鱼在药物筛选中的应用. *中国中药杂志*, 2015, 40: 1235–1239]

- 19 Tan W P, Lu H Q, Wang X T, et al. Application of zebrafish model in virus research (in Chinese). *Med Recapitul*, 2020, 26: 1673–1678 [谈文盼, 卢欢倩, 王晓桐, 等. 斑马鱼模型在病毒研究中的应用. *医学综述*, 2020, 26: 1673–1678]
- 20 Richardson R, Slanchev K, Kraus C, et al. Adult zebrafish as a model system for cutaneous wound-healing research. *J Invest Dermatol*, 2013, 133: 1655–1665
- 21 Mathias J R, Dodd M E, Walters K B, et al. Characterization of zebrafish larval inflammatory macrophages. *Dev Comp Immunol*, 2009, 33: 1212–1217
- 22 Renshaw S A, Loynes C A, Elworthy S, et al. Modeling inflammation in the zebrafish: How a fish can help us understand lung disease. *Exp Lung Res*, 2007, 33: 549–554
- 23 Progatzy F, Cook H T, Lamb J R, et al. Mucosal inflammation at the respiratory interface: A zebrafish model. *Am J Physiol-Lung Cell Mol Physiol*, 2016, 310: L551–L561
- 24 Liu L, Fokkink R, Koelmans A A. Sorption of polycyclic aromatic hydrocarbons to polystyrene nanoplastic. *Environ Toxicol Chem*, 2016, 35: 1650–1655
- 25 Boshtam M, Asgary S, Kouhpayeh S, et al. Aptamers against pro- and anti-inflammatory cytokines: A review. *Inflammation*, 2017, 40: 340–349
- 26 van der Aa L M, Chadzinska M, Tijhaar E, et al. CXCL8 chemokines in teleost fish: Two lineages with distinct expression profiles during early phases of inflammation. *PLoS One*, 2010, 5: e12384
- 27 van der Aa L M, Chadzinska M, Golbach L A, et al. Pro-inflammatory functions of carp CXCL8-like and CXCL8 chemokines. *Dev Comp Immunol*, 2012, 36: 741–750
- 28 Raghuvanshi S K, Su Y, Singh V, et al. The chemokine receptors CXCR1 and CXCR2 couple to distinct G protein-coupled receptor kinases to mediate and regulate leukocyte functions. *J Immunol*, 2012, 189: 2824–2832
- 29 Viola A, Luster A D. Chemokines and their receptors: Drug targets in immunity and inflammation. *Annu Rev Pharmacol Toxicol*, 2008, 48: 171–197
- 30 Harada A, Sekido N, Akahoshi T, et al. Essential involvement of interleukin-8 (IL-8) in acute inflammation. *J Leukoc Biol*, 1994, 56: 559–564
- 31 Kobayashi Y. The role of chemokines in neutrophil biology. *Front Biosci*, 2008, 13: 2400–2407
- 32 Tanaka T, Narazaki M, Kishimoto T. IL-6 in inflammation, immunity, and disease. *Cold Spring Harb Perspect Biol*, 2014, 6: a016295
- 33 Barton S J, Ngo S, Costello P, et al. DNA methylation of Th2 lineage determination genes at birth is associated with allergic outcomes in childhood. *Clin Exp Allergy*, 2017, 47: 1599–1608
- 34 Dris R, Gasperi J, Mirande C, et al. A first overview of textile fibers, including microplastics, in indoor and outdoor environments. *Environ Pollut*, 2017, 221: 453–458

Summary for “不同粒径聚苯乙烯微塑料和苯并(a)芘联合暴露影响炎症因子的表达”

Co-exposure to different sized polystyrene microplastics and benzo[a]pyrene affected inflammation in zebrafish and bronchial-associated cells

Yawen Xu, Jun Zhu, Jinye Hu, Zhan Zhang, Lei Li & Qian Wu*

Department of Health Inspection and Quarantine, School of Public Health, Nanjing Medical University, Nanjing 211166, China

* Corresponding author, E-mail: wuqian@njmu.edu.cn

In recent years, emerging research has revealed that microplastics (MPs) can pollute the air. Cellulose and plastic fibers have been found in the lungs of patients with lung cancer, indicating that MPs can “travel” through the atmosphere into the respiratory tract. Due to the hydrophobicity and relatively large surface area of MPs, the biotoxicity of MPs is greatly enhanced by their adsorption and enrichment in persistent organic pollutants (POPs). With the rapid development of the economy, emissions from coal burning and automobile exhaust have made polycyclic aromatic hydrocarbon (PAHs) one of the key pollutants in urban air that cannot be ignored. Epidemiological studies have shown that PAH exposure is associated with the development of respiratory diseases, such as bronchitis and asthma, in children. The aim of this study was to investigate the inflammatory effects of co-exposure to different size polystyrene MPs (PS-MPs) and benzo[a]pyrene (BaP) on adult zebrafish gills and bronchial-related cells to provide basic information for assessment of the potential health risks of combined exposure to the human respiratory system. PS-MPs (with size of 0.5 and 5 μm) and BaP were co-administered to embryos for 7 d, adult zebrafish for 14 d, and HBE cells and HBSCs for 48 h. The relative expression levels of inflammatory cytokine genes in the different exposure groups were determined by real-time polymerase chain reaction (PCR). The ingestion of microplastics into fish bodies and cells was observed by stereoscope and confocal microscopy, respectively. Result shows that microplastics of two sizes could be ingested by zebrafish embryos. When exposed only to 0.5- μm PS-MPs, relative expression of the *TNF- α* , *IL-6*, and *IL-8* genes was significantly increased compared with that in the control; relative expression of the *IFN- γ* gene was largely increased in the gills of zebrafish in 5- μm PS-MP-exposed group. Moreover, relative expression of the *IFN- γ* , *IL-6* and *IL-8* genes in the group co-exposed to BaP and 0.5- μm PS-MPs was higher than that in BaP group. In addition, 0.5- μm PS-MPs could be ingested by both cell lines. Upon exposure to 0.5- μm PS-MPs alone, relative expression of genes *TNF- α* , *IL-6*, and *IL-8* was considerably increased compared with that in the control HBE cells. In HBSCs, upon exposure to 0.5- μm PS-MPs alone, relative expression of the *TNF- α* , *IL-8*, and *IL-1 β* genes was substantially increased compared with that in the control group. Compared with that following exposure to BaP alone, relative expression of the *IFN- γ* and *IL-6* genes in HBSCs in 0.5- μm PS-MPs and BaP co-exposure group was remarkably increased, indicating a possible combined effect; however, co-exposure of 5- μm PS-MPs and BaP weakened the proinflammatory function of BaP and especially influenced relative expression of the *TNF- α* and *IL-8* genes. In addition, none of the co-exposure groups of HBE cells showed obvious combined effects. The corollary is that particle size may affect the ingestion of PS-MPs in cells. In addition, the size-dependent effects of PS-MPs influence the accumulation of organic pollutants and their biotoxicity.

polystyrene microplastics, benzo(a)pyrene, inflammation, zebrafish, cell

doi: [10.1360/TB-2020-0668](https://doi.org/10.1360/TB-2020-0668)