

# A review: Health promoting lactic acid bacteria in traditional Indonesian fermented foods

Lilis Nuraida\*

*Southeast Asian Food and Agricultural Science and Technology (SEAFST) Center, and Department of Food Science and Technology,  
Bogor Agricultural University, Indonesia*

Received 23 November 2014; received in revised form 8 December 2014; accepted 29 January 2015

## Abstract

Traditional Indonesian fermented foods can be used as potential sources of probiotics as they commonly contain lactic acid bacteria (LAB), including species of *Lactobacillus*, *Pediococcus*, *Enterococcus*, *Weisella* and *Leuconostoc*. The occurrence of LAB in Indonesian fermented foods is not only limited to lactic fermented foods but is also present in foods with molds as the main starter culture. This review aims to describe the significance of Indonesian fermented foods as potential sources of probiotics and the potential of LAB from fermented foods to promote beneficial health effects. A number of *in vitro* studies have been carried out to assess the probiotic potential of LAB from fermented foods. Many LAB strains have met the basic requirements for them to be considered as probiotics and possess some functional properties contributing to positive health impacts. Hypocholesterolemic effects, stimulation of the immune system, and prevention of diarrhea by some probiotic strains have been shown in animal studies. However, human studies on the efficacy of probiotic strains are still limited. Two strains isolated from dadih, a fermented buffalo milk, are examples of promising probiotic strains that have gone through human studies. The potential probiotic properties of LAB in Indonesian fermented foods still need to be fully investigated to assess their impact on human health. The studies should also consider factors that may influence the functional properties of probiotics, both in foods and in humans.

© 2015 Beijing Academy of Food Sciences. Production and hosting by Elsevier B.V. All rights reserved.

**Keywords:** Fermented foods; Probiotics; Diarrhea; Hypocholesterolemic; Antimutagenicity

## 1. Introduction

Fermentation is known as one of the oldest forms of food preservation in the world. Fermentation can increase the shelf-life of meat, fish, fruit and vegetables that are highly perishable due to their high water contents and nutritive values, especially in tropical countries like Indonesia. Preservation of foods occurs through lactic acid, alcoholic, acetic acid and high salt fermentations. Beside preserving foods, fermentation also changes the organoleptic characteristics of foods

through developing a wide diversity of flavors, aromas and textures. Moreover, fermentation may improve digestibility and nutritional quality through enrichment of food substrates with vitamins, proteins, essential amino acids and essential fatty acids [1,2].

As in other parts of East Asia, Indonesian fermented foods feature the use of a variety of raw materials, including cereals, soybeans, fruits, vegetables, tubers and fish. In some parts of Indonesia, meat and milk, especially buffalo milk and mare milk, have been used traditionally as raw materials for fermented products. In terms of the fermentation processes, Indonesian fermented foods can be classified into lactic fermentations (fruits, vegetables, cassava, meat, milk), alcoholic fermentations (rice, cassava), mold fermentations (soybeans, peanut press cake) and high salt fermentations (fish, soy sauce, tauco [fermented soy-bean slurry]). In the fermentation of some products, such as soy

\* Correspondence to: SEAFST Building, Jl. Puspa No. 1, IPB Darmaga Campus, Bogor, Indonesia. Tel.: +62 251 8629903; fax: +62 251 8629535.

E-mail address: [lilis@seafast.org](mailto:lilis@seafast.org)

Peer review under responsibility of Beijing Academy of Food Sciences.

sauce, a mold fermentation is followed by a brine fermentation in which LAB and yeasts are involved [3].

Although some fermentations, such as those for tempe (mold fermented soybean) and tape (alcoholic fermented steamed glutinous rice or cassava), use a starter culture, microorganisms from the environment may contaminate the ferments and grow during the fermentations. Involvement of microorganisms other than molds in tempe fermentation has started since soaking step and continues during mold fermentation [4,5]. The presence of other microorganisms such as LAB in tape fermentation contributes to the development of flavor of tape [6]. Many Indonesian fermented foods (fruits, vegetables, meat and fish) are produced through natural fermentation by controlling the environment with the addition of salt, or by soaking the raw materials in water, as in the fermentation of raw peeled cassava root. The main role of salt in fruit and vegetable fermentations is to promote the growth of LAB over spoilage bacteria [7,8] and to inhibit pectinolytic and proteolytic enzymes that can cause softening and putrefaction [7].

LAB is a group of Gram-positive, non-spore forming, coccus or rod shaped bacteria. They ferment carbohydrates to almost entirely lactic acid (homofermentation) or to a mixture of lactic acid, carbon dioxide and acetic acid and/or ethanol (heterofermentation). Other compounds, such as diacetyl, acetaldehyde and hydrogen peroxide, are also produced. These compounds contribute to the flavor and texture of fermented foods and may also contribute to the inhibition of undesirable microbes.

The LAB in Asian traditional fermented foods include *Lactobacillus plantarum*, *Lb. pentosus*, *Lb. brevis*, *Lb. fermentum*, *Lb. casei*, *Leuconostoc mesenteroides*, *Leu. kimchi*, *Leu. fallax*, *Weissella confusa*, *W. koreensis*, *W. cibaria*, and *Pediococcus pentosaceus*, many of which are considered to be potential probiotics [7]. Most of the LAB present in Indonesian fermented foods are *Lactobacillus* species (Table 1). Other genera, such as *Pediococcus*, *Lactococcus*, *Enterococcus*, *Weissella* and *Leuconostoc*, are also found in some fermented foods (Table 1). LAB are involved to varying degrees in Asian fermented foods, and may have positive and negative effects on products [9]. In cereal alcoholic fermentations, lactic acid bacteria contribute to the characteristic of flavor and taste. Excessive lactic acid generally lowers the quality of alcoholic fermentation products. However, in fruit, vegetable, milk and meat fermentations, LAB play a major role in producing acid necessary to the quality of the products. It is interesting that LAB are generally present in tempe, which is not an acidic fermentation. In tempe fermentation, soybeans are soaked overnight prior to inoculation with starter culture containing *Rhizopus oligosporus* as the primary microorganism. Acid fermentation involving LAB takes place during the soaking [10,11] and some growth of lactic acid bacteria commonly occurs during the stage of mold growth [5,12].

LAB in fermented foods are of interest not only for their role in fermentation but also for their role in promoting positive health impacts. The concept of beneficial health effects of LAB has existed since Metchnikoff in 1908 proposed that

acid producing microorganisms in fermented dairy products could lead to a prolongation of the life span of consumers [36]. Although historically the fermented products associated with beneficial LAB were milk-based, much research has been directed to exploring LAB from other fermented foods as potential probiotics. A probiotic is defined as a live microorganism that will confer beneficial effects on the host when ingested in sufficient amount [37]. The probiotic bacteria used in commercial products are mainly members of the genera *Lactobacillus* and *Bifidobacterium* [36]. Probiotic bacteria are usually those bacteria that have adapted to the gastrointestinal environment. However, recent research has shown promising probiotic activity of LAB isolated from fermented foods [9]. The progress in research on the beneficial health effects of microorganisms, especially LAB, isolated from Indonesian fermented foods is discussed below.

## 2. Potential of lactic acid bacteria isolated from Indonesian fermented foods as probiotics

Probiotic and other functional properties are strain dependent and all probiotic strains are unique and different; therefore, their properties and characteristics need to be well defined [38]. Several criteria have to be met in selecting probiotic strains, including acid and bile tolerance, survival through the gastrointestinal track, ability to adhere to intestinal surfaces, antimicrobial activity against potentially pathogenic bacteria, and good technological properties [39]. Functional properties of probiotics include hypocholesterolemic activity by lowering plasma cholesterol [40], preventing and treating diarrhea [37], and altering the immune system [41,42]. The mechanisms by which probiotics exert their beneficial effects on the host include the reduction of luminal pH, competition with pathogens for adhesion sites and nutritional sources, secretion of antimicrobial substances, toxin inactivation, and immune stimulation [43].

Based on *in vitro* studies, LAB isolated from Indonesian fermented foods have promising characteristics as probiotic candidates (Table 2). *In vitro* assessment shows that many LAB isolates tolerate bile salt and low pH environment and possess antagonistic activity against foodborne pathogens. These characteristics are similar to those of intestinal microorganisms, such as *Lactobacillus acidophilus* and *Lb. casei* that are commonly used as probiotics [9]. The research results suggest that the LAB in Indonesian fermented food have adapted to environments that resemble the gastrointestinal track and, hence, have potential as probiotic microorganisms. Adaptation of LAB isolated from fermented foods on specific environment such as high salt concentration, acidic condition, has also been reported such as *L. mesenteroides* in Kimchii, cane juice, *Leu. oenos* in grape juice and *Tetragenococcus halophilus* in soy sauce [9]. It has been suggested that adaptation involves the human food cycle, from soil to raw materials, to fermented product, to human intestine, to feces and then to soil again [9].

Table 1  
Occurrence of lactic acid bacteria in some Indonesian fermented foods.

| Fermented food                        | Main raw material(s)                                                             | Fermentation process                                                      | Lactic acid bacteria present                                                                                                                                                                                                                                                                                                                                                                                                                                      | References                              |
|---------------------------------------|----------------------------------------------------------------------------------|---------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------|
| Sayur asin                            | Mustard cabbage leaf                                                             | Lactic fermentation                                                       | <i>Lactobacillus farciminis</i> , <i>Lb. fermentum</i> ,<br><i>Lb. namurensis</i> , <i>Lb. plantarum</i> , <i>Lb.</i><br><i>helveticus</i> , <i>Lb. brevis</i> , <i>Lb. versmoldensis</i> ,<br><i>Lb. casei</i> , <i>Lb. rhamnosus</i> , <i>Lb.</i><br><i>fabifermentans</i> , <i>Lb. satsumensis</i><br><i>Leuconostoc mesenteroides</i> , <i>Lb.</i><br><i>confusus</i> , <i>Lb. curvatus</i> , <i>Pediococcus</i><br><i>pentosaceus</i> , <i>Lb. plantarum</i> | [13]<br>[7]                             |
| Tempoyak                              | Flesh of durian ( <i>Durio zibethinus</i> )                                      | Lactic fermentation                                                       | <i>Lb. plantarum</i> , <i>Lb. coryneformis</i> , <i>Lb.</i><br><i>casei</i><br><i>Lb. plantarum</i> , <i>Lactobacillus</i> sp.,<br><i>Weissella paramesenteroides</i> ,<br><i>Pediococcus acidilactici</i><br><i>Enterococcus gallinarum</i> , <i>E. faecalis</i>                                                                                                                                                                                                 | [14]<br>[15]<br>[16]                    |
| Mandai                                | Flesh of cempedak, a family of jack fruit ( <i>Artocarpus champeden</i> Spreng.) | Lactic fermentation                                                       | <i>Lb. plantarum</i><br><i>P. pentosaceus</i>                                                                                                                                                                                                                                                                                                                                                                                                                     | [17,18]<br>[18]                         |
| Tape starter culture                  | Rice flour                                                                       | –                                                                         | <i>P. pentosaceus</i> , <i>E. faecium</i> , <i>Lb. curvatus</i> ,<br><i>W. confusa</i> , <i>W. paramesenteroides</i>                                                                                                                                                                                                                                                                                                                                              | [19]                                    |
| Rice wine/rice tape                   | Glutinous rice (steamed)                                                         | Alcoholic fermentation                                                    | <i>P. pentosaceus</i> , <i>Weissella</i> sp.                                                                                                                                                                                                                                                                                                                                                                                                                      | [20]                                    |
| Growol                                | Cassava (raw)                                                                    | Lactic fermentation                                                       | <i>Lb. plantarum</i> , <i>Lb. rhamnosus</i>                                                                                                                                                                                                                                                                                                                                                                                                                       | [21,22]                                 |
| Tempe                                 | Soybean                                                                          | Mold fermentation                                                         | <i>Lb. fermentum</i> , <i>Lb. plantarum</i> , <i>P.</i><br><i>pentosaceus</i> , <i>W. confusa</i> , <i>Lb. delbrueckii</i><br><i>ssp. delbrueckii</i><br><i>Lb. plantarum</i>                                                                                                                                                                                                                                                                                     | [23]<br>[24]                            |
| Soy sauce (kecap)                     | Soybean                                                                          | Mold fermentation followed by high salt fermentation (brine fermentation) | <i>Tetragenococcus halophilus</i>                                                                                                                                                                                                                                                                                                                                                                                                                                 | [3]                                     |
| Bakasang                              | Fish                                                                             | High salt fermentation                                                    | <i>P. acidilactici</i>                                                                                                                                                                                                                                                                                                                                                                                                                                            | [25]                                    |
| Urutan (traditional Balinese sausage) | Lean Pork                                                                        | Lactic fermentation                                                       | <i>Lb. plantarum</i> , <i>P. acidilactici</i> , <i>Lb.</i><br><i>faricimins</i>                                                                                                                                                                                                                                                                                                                                                                                   | [26]                                    |
| Dadih                                 | Buffalo milk                                                                     | Lactic fermentation                                                       | <i>Leu. mesenteroides</i><br><i>Lactococcus lactis</i> subsp. <i>lactis</i> , <i>Lb.</i><br><i>brevis</i> , <i>Leu. mesenteroides</i> , <i>Lb. casei</i><br><i>Lb. plantarum</i> , <i>E. faecium</i><br><i>Lb. fermentum</i> , <i>Leu. lactis</i> subsp. <i>lactis</i><br><i>Lc. lactis</i> , <i>Lb. rhamnosus</i>                                                                                                                                                | [27]<br>[28]<br>[29,30]<br>[31]<br>[32] |
| Fermented mare milk                   | Mare milk                                                                        | Lactic fermentation                                                       | <i>Lb. rhamnosus</i> , <i>Lb. fermentum</i><br><i>Lb. acidophilus</i> , <i>Lb. brevis</i>                                                                                                                                                                                                                                                                                                                                                                         | [33,34]<br>[35]                         |

Adhesion properties similar to those of probiotic strains have also been shown by some isolates from fermented foods, such as *Lactobacillus reuteri* IS-27560, *Lactococcus lactis* IS-16183 and *Lb. rhamnosus* IS-7257 from dadih that adhere to mucus layers and Caco-2 cells [32]. *Lc. lactis* IS-16183 and *Lb. rhamnosus* IS-7257 significantly inhibited adhesion of *Escherichia coli* O157:H7. Accordingly, these two strains may be potential candidates for use as probiotic strains. The adhesion properties of dadih isolates were relatively comparable to the commercial probiotic strains, *Lb. casei* Shiota and *Lb. rhamnosus* GG. Another study on dadih isolates (*Lb. plantarum* IS-10506 and IS-20506; *Enterococcus faecium* IS-27526, IS-23427 and IS-16183) showed that *Lb. plantarum* IS-10506 was the most adhesive and can significantly reduce

pathogen adhesion to mucus [29]. *Lb. plantarum*, isolated from fermented fruit (mandai), also showed good adhesive properties on enterocyte-like HCT-116 cells, with *Lb. plantarum* MB427 showing the strongest inhibition of adhesion of *Listeria monocytogenes* ATCC 13932, enteropathogenic *E. coli* (EPEC) K1.1 and *Salmonella enterica* serovar Typhimurium ATCC 14028 [17]. These findings suggest that antimicrobial effects of fermented food isolates also involve adhesion properties. Production of substances other than organic acids as shown by certain isolates from bakasang, dadih and growol [25,28,44] may enhance their activity in inhibiting the growth of pathogens.

Among potential health beneficial effects, *in vitro* assessment of fermented foods isolates for their hypocholesterolemic

Table 2

Characteristic and functional properties of lactic acid bacteria isolated from Indonesian fermented foods based on *in vitro* studies.

| Lactic acid bacteria                                                                                                                                                                                                          | Source of LAB                                          | Characteristic and functional properties                                                                                                                                                                                                                                                                 | References |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| <i>Enterococcus gallinarum</i> UP-9, <i>Enterococcus faecalis</i> UP-11                                                                                                                                                       | Tempoyak                                               | Bile and acid tolerant, being able to conjugate sodium taurocholate                                                                                                                                                                                                                                      | [16]       |
| <i>Lb. plantarum</i>                                                                                                                                                                                                          | Mandai                                                 | Bile salt and low pH tolerant, good adhesion properties                                                                                                                                                                                                                                                  | [17]       |
| <i>Lactobacillus plantarum</i> sa28k                                                                                                                                                                                          | Indonesian sauerkraut                                  | Acid and bile resistant, assimilation of cholesterol                                                                                                                                                                                                                                                     | [45]       |
| <i>Lb. acidophilus</i> FNCC116                                                                                                                                                                                                | Moromi soy sauce fermentation                          |                                                                                                                                                                                                                                                                                                          |            |
| <i>Lb. casei</i> FNCC262                                                                                                                                                                                                      | Tape ketan                                             |                                                                                                                                                                                                                                                                                                          |            |
| <i>Lb. fermentum</i>                                                                                                                                                                                                          | Tempe                                                  | Bile salt and low pH tolerant                                                                                                                                                                                                                                                                            | [23]       |
| <i>Lactobacillus</i> TGR-2                                                                                                                                                                                                    | Growol                                                 | Inhibits growth of <i>Staphylococcus aureus</i> , <i>S. typhimurium</i> , <i>E. coli</i> , <i>Bacillus cereus</i> , <i>Morganella morganii</i> . Produces bacteriocin-like compounds                                                                                                                     | [44]       |
| <i>Lb. casei</i> FNCC262                                                                                                                                                                                                      | Tape ketan (alcoholic fermented glutinous rice)        | Tolerance to 0.3% bile salt, resistant to pH 2.5, antimicrobial activity against <i>E. coli</i> , <i>S. aureus</i> , <i>B. cereus</i>                                                                                                                                                                    | [51]       |
| <i>Lb. acidophilus</i> FNCC 116                                                                                                                                                                                               | Moromi soy sauce (brine fermentation)                  |                                                                                                                                                                                                                                                                                                          |            |
| DA-1                                                                                                                                                                                                                          | Dadih                                                  |                                                                                                                                                                                                                                                                                                          |            |
| DA-2                                                                                                                                                                                                                          | Dadih                                                  |                                                                                                                                                                                                                                                                                                          |            |
| <i>Lactococcus lactis</i> subsp. <i>lactis</i> IS-10285, IS-7386, IS-16183, IS-11857 and IS-29862, <i>Lb. brevis</i> IS-27560, IS-26958 and IS-23427, <i>Leuconostoc mesenteroides</i> IS-27526, and <i>Lb. casei</i> IS-7257 | Dadih                                                  | Bile salt (oxgall), low pH and lysozyme tolerant                                                                                                                                                                                                                                                         | [28]       |
| <i>Lb. plantarum</i> Dad-3                                                                                                                                                                                                    | Dadih                                                  | <i>Lc. lactis</i> subsp. <i>lactis</i> IS-11857 and IS-29862, <i>Lb. brevis</i> IS-26958 showed high alt hydrolase (BSH) activity. <i>Lc. lactis</i> subsp. <i>lactis</i> IS-10285 and IS-16183 had a positive spectrum of bacteriocin activity against <i>E. coli</i> and <i>Listeria monocytogenes</i> |            |
| <i>Lb. plantarum</i> Mut 7 and Mut 13                                                                                                                                                                                         | Gatot (fermented cassava)                              | Inhibiting <i>Shigella dysenteriae</i> , <i>E. coli</i> and <i>S. typhi</i>                                                                                                                                                                                                                              | [52]       |
| <i>Lb. plantarum</i> T-3                                                                                                                                                                                                      | Growol (fermented cassava)                             | Inhibiting <i>Shigella dysenteriae</i>                                                                                                                                                                                                                                                                   |            |
| <i>Lb. fermentum</i> I-11 and <i>Leu. lactis</i> subsp. <i>lactis</i> I-2775                                                                                                                                                  | Dadih                                                  | Acid and oxgall (bile) tolerant, deconjugated sodium taurocholate and bound cholesterol                                                                                                                                                                                                                  | [31]       |
| <i>Lb. reuteri</i> IS-27560, <i>Lc. lactis</i> IS-16183, <i>Lb. rhamnosus</i> IS-7257, <i>Enterococcus faecium</i> IS-27526                                                                                                   | Dadih                                                  | Adherence to mucus layer and Caco-2 cells. IS 16183 and IS7257 inhibit the adhesion of <i>Escherichia coli</i> O157:H7 to human intestinal mucosal surface                                                                                                                                               | [32]       |
| <i>Lb. plantarum</i> IS-10506, IS-20506 and <i>E. faecium</i> IS-27526, IS-23427 and IS-16183                                                                                                                                 | Dadih                                                  | Adhesion was strain dependent with the most adhesive was <i>Lb. plantarum</i> strain was IS-10506 and reduced pathogen adhesion to mucus                                                                                                                                                                 | [29]       |
| <i>Lb. plantarum</i> IS-10507 and IS-20506                                                                                                                                                                                    | Dadih                                                  | Removal of microcystin-LR, a cyclic heptapeptide hepatotoxin (cyanobacterial toxin)                                                                                                                                                                                                                      | [47]       |
| <i>Lb. brevis</i> , <i>Lb. acidophilus</i>                                                                                                                                                                                    | Fermented mare milk                                    | Survive in low pH (2.5) for 2 h, bile tolerant                                                                                                                                                                                                                                                           | [35]       |
| <i>Lb. rhamnosus</i> FSMM15, FSMM22, FSMM26                                                                                                                                                                                   | Fermented mare milk                                    | Bile salt, low pH and artificial gastrointestinal fluids tolerant; and having good adhesion properties                                                                                                                                                                                                   | [33]       |
| <i>P. acidilactici</i>                                                                                                                                                                                                        | Bekasam/bekasang                                       | Antimicrobial activity against <i>E. coli</i> , <i>S. aureus</i> , <i>P. fluorescens</i>                                                                                                                                                                                                                 | [24]       |
| LAB (unidentified)                                                                                                                                                                                                            | Bekasam                                                | Antimicrobial activity against <i>E. coli</i> , <i>S. typhimurium</i> , <i>B. cereus</i> , <i>S. aureus</i>                                                                                                                                                                                              | [53]       |
| <i>Lb. plantarum</i> SMN 025, <i>Lb. casei</i> subsp. <i>rhamnosus</i> FNCC 098                                                                                                                                               | Fermented foods                                        | $\beta$ -Glucosidase activity                                                                                                                                                                                                                                                                            | [48]       |
| <i>Pediococcus acidilactici</i> , <i>P. pentosaceus</i> , <i>P. lolii</i> AB362985, <i>Lb. pentosus-plantarum</i> group                                                                                                       | Fermented foods (tape, tempe and fermented vegetables) | $\beta$ -Glucosidase activity                                                                                                                                                                                                                                                                            | [49]       |

effect has been done by some researchers [16,31,45]. Cholesterol binding by LAB in the small intestine may reduce the amount of dietary cholesterol absorbed [31]. Several mechanisms for lowering cholesterol absorption have been hypothesized [46], including enzymatic deconjugation of bile acids by bile-salt hydrolase (BSH), assimilation of cholesterol, co-precipitation of cholesterol with deconjugated bile, binding of cholesterol to the cell walls of probiotic bacteria, incorporation of cholesterol into the cell membranes of probiotics, conversion of cholesterol to coprostanol, and production of short-chain fatty acids by probiotics in the presence of prebiotic substrates. BSH hydrolyzes conjugated glycodeoxycholic acid and taurodeoxycholic acid, leading to the deconjugation of glyco- and tauro-bile acids. Once deconjugated, bile acids are less soluble and absorbed by the intestines, leading to their elimination in the feces. Cholesterol is used to synthesize new bile acids in a homeostatic response, resulting in lowering of serum cholesterol. The ability of cholesterol-binding appeared to be growth and strain specific.

Other potential health benefits shown by LAB isolates from fermented foods are the ability of a *L. plantarum* isolate from dadih to remove microcystin-LR, a cyclic heptapeptide hepatotoxin produced by cyanobacteria [47] and the ability to produce  $\beta$ -glucosidase [48,49]. Cyanobacteria produce a number of potent hepato- and neurotoxins, collectively called cyanotoxins, which have potent acute hepatotoxicity and tumor promoting activity [47]. *Lb. plantarum* IS-10506 and IS-20506 have shown ability to efficiently remove the toxins. Meanwhile,  $\beta$ -glucosidase activity is widespread among LAB and presumably plays a role in interactions with the human host [50].  $\beta$ -Glucosidase releases a wide range of plant secondary metabolites from their  $\beta$ -D-glucosylated precursors. The conversion of glucoside isoflavones into their bioactive aglycones by LAB has been observed in soymilk fermentation [48].

LAB isolated from fermented foods has potential as producers of bioactive compounds. Gamma-aminobutyric acid (GABA) has various physiological functions and could be produced by LAB [54]. Some *Lactobacillus* species have been reported to produce equol [7-hydroxy-3-(40-hydroxyphenyl) chroman], a nonsteroidal estrogen of the isoflavone class in fermented soymilk [55].

Most established probiotics are LAB and Bifidobacteria although recently certain yeasts (e.g. *Saccharomyces boulardii* [56–58]) and spore-forming bacteria (e.g. *Bacillus coagulans* [59–61]) have been considered as probiotics. While *S. boulardii* has been shown to be effective in preventing the recurrence of *Clostridium difficile*-induced pseudomembranous colitis as well as the antagonistic action of *E. coli*, the long-term advantages of using spores as probiotics is that they are heat-stable and can survive transit across the stomach barrier, properties that cannot be assured with other probiotic bacteria that are given in the vegetative form [61]. Yeasts are commonly present in many fermented foods [6,11,12,62] and starter culture [63]. Spore-forming bacteria could also be part of the microbial consortium in traditional Indonesian fermented foods, such as tempe [5] and tape [6,64] as well as their starter cultures [62].

### 3. Health beneficial effect of lactic acid bacteria isolated from fermented foods

#### 3.1. Animal studies

Only a few promising LAB isolated from Indonesian fermented foods have been assessed in animal studies for their potential health benefits (Table 3). Evaluation of *L. plantarum* sa28k, *Lb. acidophilus* FNCC116 and *Lb. casei* FNCC262 in lowering cholesterol in rats, revealed that rats that received milk fermented by the three LAB had significantly lower serum cholesterol levels than rats feeds non-fermented milk [45]. *Lb. acidophilus* KBc and *Lb. brevis* KBa, from fermented mare milk, were able to adhere and colonize gut mucosal epithelium of rats [35]. Administration of these isolates significantly reduced cholesterol levels in blood serum of hypercholesterolemic rabbits. The cholesterol-lowering activity of milk fermented by *Lactococcus lactis* subsp. *lactis* IS-10285 and *Lc. lactis* subsp. *lactis* IS-29862 has been evaluated in hypercholesterolemic rats [65]. The isolates had high taurocholate-deconjugating activity. Only milk fermented by *Lc. lactis* subsp. *lactis* IS-10285 significantly reduced the total serum cholesterol, LDL cholesterol and total bile acids. Neither milk nor fermented milk influenced HDL cholesterol levels. The authors suggested that the hypocholesterolemic effect of *Lc. lactis* subsp. *lactis* IS-10285 was due to its ability to suppress the reabsorption of bile acids into the enterohepatic circulation and enhance the excretion of bile acids in feces of hypercholesterolemic rats. These results indicate that strains of LAB isolated from Indonesian fermented foods could be considered as probiotic strains that have beneficial effects in reducing serum cholesterol levels.

Candidate probiotic strains isolated from Indonesian fermented foods have also been evaluated in the treatment or prevention of diarrhea caused by Enteropathogenic *E. coli* (EPEC) infection. *Lb. plantarum* MB427 from mandai at  $10^9$  cfu/mL reduced the incidence and severity of diarrhea and shortened the duration of diarrhea in EPEC-induced diarrhea in Sprague-Dawley rats [17] and increased secretion of serum IgA and IgB. Supplementation of dadih isolate, *P. pentosaceus*, at a dose of  $2 \times 10^8$  cfu/g reduced stool frequency, lowered tumor necrosis factor- $\alpha$  levels and improved the balance of gut microflora in EPEC-induced diarrheal mice (*Mus musculus*) [66]. Similar results were observed with a *W. paramesenteroides* strain also isolated from dadih [67]. The researchers suggested that probiotic supplementation may protect against mucosal epithelial cell damage by *E. coli* exposure and protect cell against further damage by TNF- $\alpha$  and interferon (IFN)- $\gamma$ . Probiotics are able to downregulate T helper (Th)-1 responses and inhibit the production of proinflammatory cytokines, such as TNF- $\alpha$  by dendritic cells. The decreased the concentration of TNF- $\alpha$  in the feces, decreased the serum levels of TNF- $\alpha$ , and stool frequency. These studies indicate that potential probiotic bacteria isolated from fermented foods could improve the immune system and reduce diarrhea incidents. Diarrhea is the third leading cause of death, following tuberculosis and pneumonia, in Indonesia [66].

Table 3

Health benefits of lactic acid bacteria isolated from Indonesian fermented foods as assessed in animal studies.

| Lactic acid bacteria                                                                               | Source of LAB                                                        | Characteristic and functional properties                                                                                                       | References |
|----------------------------------------------------------------------------------------------------|----------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| <i>Lb. plantarum</i> MB 427                                                                        | Mandai                                                               | Shortened duration of diarrhea in rats caused by EPEC infection and induced secretion of IgA and IgG                                           | [17]       |
| <i>Lactobacillus plantarum</i> sa28k<br><i>Lb. acidophilus</i> FNCC116<br><i>Lb. casei</i> FNCC262 | Indonesian sauerkraut<br>Moromi soy sauce fermentation<br>Tape ketan | Assimilating cholesterol and lowering serum cholesterol in rats                                                                                | [45]       |
| <i>Lb. acidophilus</i> KBc and <i>Lb. brevis</i> Kba                                               | Fermented mare milk                                                  | Adhered to and colonized gut mucosa epithelium of rat, reduced cholesterol level of blood serum of rabbits with hypercholesterolemia condition | [35]       |
| <i>Lactococcus lactis</i> subsp. <i>lactis</i> IS-10285                                            | Dadih                                                                | Significantly reduced serum total cholesterol, LDL cholesterol and total bile acids in hypercholesterolemic rats                               | [65]       |
| <i>P. pentosaceus</i>                                                                              | Dadih                                                                | Reduced stool frequency, lowered TNF- $\alpha$ level and improved the balance of gut microflora in EPEC-induced diarrheal mice                 | [66]       |
| <i>W. mesenteroides</i>                                                                            | Dadih                                                                | Reduce stool frequency, lower TNF- $\alpha$ level and improve the balance of gut microflora in EPEC-induced diarrhea mice                      | [67]       |
| <i>Enterococcus faecium</i> IS-27526                                                               | Dadih                                                                | Significantly lowered fecal mutagenicity of rats fed with milk cultured with the isolate                                                       | [30]       |

The most common causes of diarrhea in children, both in developed and developing countries are *E. coli*, Rotavirus, *Salmonella* spp., *Shigella* spp., *Campylobacter jejuni*, *Entamoeba histolytica* and *Giardia lamblia* [68]. The promising features of LAB isolated from fermented foods in reducing/preventing the incident of diarrhea needs to be confirmed in human studies.

Modification of gut bacterial activities has also correlated with antimutagenicity. Potential probiotic bacterium, *E. faecium* IS-27526 isolated from dadih, showed *in vivo* anti-mutagenic properties toward Trp-P1 of rats [30]. Milk cultured with *E. faecium* IS-27526 significantly lowered fecal mutagenicity and the recovery of Trp-P1 in urine was significantly lower than in control rats fed skim milk. The anti-mutagenic properties was considered due to binding ability of the bacterial cell wall of dadih lactic cultures toward chemicals mutagen.

### 3.2. Human studies

Human studies on LAB isolated from Indonesian fermented foods are very limited. Two potential probiotics originating from dadih have gone through a clinical study in humans [69,70]. Surono et al. [69] evaluated the effect of *E. faecium* IS-27526 in milk on humoral immune response and on body-weight of young children aged between 15 and 54 months. A 90 days randomized, double-blind, placebo-controlled study was conducted with two groups of young children, placebo and probiotic group. Ultra high temperature treated, low fat milk was used as the carrier of  $2.3 \times 10^8$  cfu/day of the probiotic. The results showed that *E. faecium* IS-27526 had a significant positive effect on humoral immune response and salivary IgA in underweight young children, and on their weight gain. However, the total serum IgA did not significantly increase in the probiotic group compared with the placebo group.

Another human study [70] evaluated the effect of probiotic *L. plantarum* IS-10506 (originally isolated from dadih) and zinc supplementation on humoral immune response and zinc status of Indonesian infants aged 12–24 months in a 90-day randomized, double-blind, placebo-controlled trial. A combination of microencapsulated *Lb. plantarum* IS-10506 at a dose of  $10^{10}$  cfu/day and 8 mg of elemental zinc showed a potential ability to improve the zinc status of the infants. Supplementation with the probiotic and zinc resulted in a significantly increased humoral immune response, as well as improved zinc status. The effects of probiotics on the humoral immune response was suggested to be the result of colonization and adhesion to epithelial cells, and production of SIgA production induced by the cell wall component of probiotics, such as lipoteichoic acids and peptidoglycan [69]. Both strong immune responses and gut integrity play important roles in the repair of intestinal brush border damage as a result of the balance of microbiota. Zinc and probiotics work *via* different mechanisms, but they are possible to have a synergistic effect. While zinc is necessary for the activity of some immunity mediators, probiotic supplementation could improve the integrity of the intestine, which in turn will optimize the absorption of zinc. The researchers considered that the health beneficial effects of zinc and probiotics could be amplified when they are taken together, more specifically, resulting in improved mineral absorption or a higher cellular immune response.

### 4. Future perspectives

Many traditional Indonesian fermented foods are potential sources of microorganisms, especially LAB, with promising beneficial health effects. Among the functional properties explored, LAB originating from Indonesian fermented foods showed promising effects on hypocholesterolemia, stimulation of the immune system, and in the prevention of diarrhea. These

beneficial health effects could contribute to promoting the health status of Indonesians. However, the functional properties of many promising LAB originating from Indonesian fermented foods have not been fully investigated. Studies are mainly of a preliminary nature and need to be confirmed in animal and human studies. Although most Indonesian fermented foods are historically safe, exploiting a single microorganism should also be supported with studies to confirm its safety, especially for strains belonging to genera that have been reported to pose safety issues. In addition, the ability of LAB of to produce bioactive compounds is an area that needs to be explored extensively in order to maximize the beneficial health impact.

Consumers are aware that fermented foods contain microorganisms and, hence, they become suitable matrixes as probiotic carriers. Among Indonesian fermented products, tempe, as a cheap meat substitute, is an important part of the daily diet. However, tempe is commonly consumed after cooking, which would kill any live probiotic LAB. Other fermented foods, such as *dadih*, *tape*, *sayur asin* and beverages, that are consumed without cooking are more suitable as probiotic carriers. When fermented foods are used as carriers of probiotic bacteria, factors that may influence the ability of the probiotic to survive in the product, during fermentation and marketing, and their activity when entering the human gastrointestinal track, must be considered. LAB with beneficial health effects in fermented foods could serve both as probiotics and as the fermentative organism. Ideally, probiotic cultures to be incorporated into fermented products should have multifunctional characteristics. They should be able to grow and ferment the product, or at least not have any negative impact on the organoleptic properties of the food, while maintaining their probiotic properties.

Exploration of potentially beneficial microorganisms in Indonesian fermented foods needs to be extended to microorganisms other than LAB that are also present in fermented foods. Yeasts and some species of *Bacillus* for examples are the potential microorganisms for probiotic.

Traditional Indonesian fermented foods, utilizing a variety of raw materials, have been handed down for many generations. Although challenges remain, the microorganisms present and involved during fermentation as well as their metabolic products may contribute health benefits to consumers. Therefore, more evidence on the health impacts of beneficial microorganisms and understanding of the relationships between fermented foods, beneficial microorganisms and human health are essential if use of beneficial microorganisms to promote health is to be fully exploited.

## Acknowledgement

The author would like to thank Dr. J.D. Owens of University of Reading, UK for his assistance in reading and correcting the manuscript.

## References

- [1] K.H. Steinkraus, Nutritional significance of fermented foods, *Food Res. Int.* 27 (3) (1994) 259–267.
- [2] K.H. Steinkraus, Fermentations in world food processing, *Compr. Rev. Food Sci. Food Saf.* 1 (2002) 24–32.
- [3] W.F.M. Roling, A. Apriyanto, H.W. Van Verseveld, Comparison between traditional and industrial soy sauce (*Kecap*) fermentation in Indonesia, *J. Ferment. Bioeng.* 81 (3) (1996) 275–278.
- [4] J.D. Owens, M. Astuti, K.R. Kuswanto, Tempe and related products, in: J.D. Owens (Ed.), *Indigenous Fermented Foods of Southeast Asia*, CRC Press, 2014, pp. 1–108.
- [5] T. Barus, A. Suwanto, A.T. Wahyudi, et al., Role of bacteria in tempe bitter taste formation and molecular biological analysis base on 16S rRNA gene, *J. Microbiol. Indones.* 2 (2008) 17–21.
- [6] L. Nuraida, J.D. Owens, Sweet, sour, alcoholic solid substrate fungal fermentations, in: J.D. Owens (Ed.), *Indigenous Fermented Foods of Southeast Asia*, CRC Press, 2014, pp. 137–155.
- [7] M.R. Swain, M. Anandharaj, R.C. Ray, et al., Fermented fruits and vegetables of Asia: a potential source of probiotics, *Biotechnol. Res. Int.* (2014) 19, Article ID 250424.
- [8] L. Nuraida, J.D. Owens, J.A. Bakar, et al., Lactic vegetable and fruit fermentations, in: J.D. Owens (Ed.), *Indigenous Fermented Foods of Southeast Asia*, CRC Press, 2014, pp. 185–209.
- [9] S.J. Rhee, J.E. Lee, C.H. Lee, Importance of lactic acid bacteria in Asian fermented foods, *Microb. Cell Fact.* 10 (1) (2011) S5.
- [10] R.K. Mulyowidarso, G.H. Fleet, K.A. Buckle, The microbial ecology of soybean soaking for tempe production, *Int. J. Food Microbiol.* 8 (1989) 35–46.
- [11] M.J.R. Nout, J.L. Kiers, Tempe fermentation, innovation and functionality: update into the third millennium, *J. Appl. Microbiol.* 98 (2005) 789–805.
- [12] Efriwati, A. Suwanto, G. Rahayu, et al., Population dynamics of yeasts and lactic acid bacteria (LAB) during tempeh production, *HAYATI J. Biosci.* 20 (2) (2013) 57–64.
- [13] Sulistiani, Abinawanto, E. Sukara, et al., Identification of lactic acid bacteria in sayur asin from Central Java (Indonesia) based on 16S rDNA sequence, *Int. Food Res. J.* 21 (2) (2014) 527–532.
- [14] C.U. Wirawati, Potency of Lactic Acid Bacteria Isolated from Tempoyak as Probiotic (Thesis in Indonesian), Bogor Agricultural University, Bogor, 2002.
- [15] N. Yuliana, E.I. Dizon, Phenotypic identification of lactic acid bacteria isolated from *Tempoyak* (fermented durian) made in the Philippines, *Int. J. Biol.* 3 (2) (2011) 145–152.
- [16] U. Pato, I.S. Surono, Bile and acid tolerance of lactic acid bacteria isolated from tempoyak and their probiotic potential, *Int. J. Agric. Technol.* 9 (7) (2013) 1849–1862.
- [17] A. Emmawati, Study of Antiinfection Properties of Lactic Acid Bacteria Isolated from Mandai (Dissertation), Graduate School, Bogor Agricultural University, Indonesia, 2014.
- [18] E.S. Rahayu, Lactic acid bacteria in fermented foods of Indonesian origins, *AgriTech* 23 (2003) 75–84.
- [19] I.N. Sujaya, S. Amachi, A. Yokota, et al., Identification and characterization of lactic acid bacteria in ragi tape, *World J. Microbiol. Biotechnol.* 17 (2001) 349–357.
- [20] I.N. Sujaya, S. Amachi, K. Saito, et al., Specific enumeration of lactic acid bacteria in ragi tape by colony hybridization with specific oligonucleotide probes, *World J. Microbiol. Biotechnol.* 18 (2002) 263–270.
- [21] W.D.W. Putri, M. Mika, Y. Nakagawa, et al., Differentiation studies of predominant lactic acid bacteria isolated during *growol* fermentation by using polyphasic taxonomic characterization, *Afr. J. Biotechnol.* 10 (42) (2011) 8194–8204.
- [22] W.D.R. Putri, H. Haryadi, D.W. Marseno, et al., Isolation and characterization of amylolytic lactic acid bacteria during growol fermentation, an Indonesian traditional food, *J. Teknol. Pert.* 13 (1) (2012) 52–60.
- [23] K. Touw, Identification of Dominant Lactic Acid Bacteria During Tempeh Fermentation and Evaluation of Their Potential as a Probiotic, Faculty of Agricultural Technology, Bogor Agricultural University, 2014 (research report).
- [24] S. Kormin, G. Rusul, S. Radu, et al., Bacteriocin-producing lactic acid bacteria isolated from traditional fermented food, *Malays. J. Med. Sci.* 8 (1) (2001) 63–68.

- [25] H.J. Lawalata, L. Sembiring, E.S. Rahayu, Molecular identification of lactic acid bacteria producing antimicrobial agents from bakasang, an Indonesian traditional fermented fish product, *Indones. J. Biotechnol.* 16 (2) (2011) 93–99.
- [26] N.S. Antara, I.N. Sujaya, A. Yokota, et al., Identification and succession of lactic acid bacteria during fermentation of 'urutan', a Balinese indigenous fermented sausage, *World J. Microbiol. Biotechnol.* 18 (2002) 255–262.
- [27] A. Hosono, R. Wardoyo, H. Otani, Microbial flora in "dadih", a traditional fermented milk in Indonesia, *Lebensm. Wiss. Technol.* 22 (1989) 20–24.
- [28] I.S. Surono, *In vitro* probiotic properties of indigenous dadih lactic acid bacteria, *Asian-Australas. J. Anim. Sci.* 16 (5) (2003) 726–731.
- [29] M.C. Collado, I.S. Surono, J. Meriluoto, et al., Potential probiotic characteristics of *Lactobacillus* and *Enterococcus* strains isolated from traditional dadih fermented milk against pathogen intestinal colonization, *J. Food Prot.* 70 (3) (2007) 700–705.
- [30] I.S. Surono, U. Pato, Koesnandar, et al., *In vivo* antimutagenicity of dadih probiotic bacteria towards Trp-P1, *Asian-Australas. J. Anim. Sci.* 22 (1) (2009) 119–123.
- [31] U. Pato, M. Ali, A.K. Parlindungan, Taurocholate deconjugation and cholesterol binding by indigenous dadih lactic acid bacteria, *Hayati* 12 (3) (2005) 103–107.
- [32] J. Dharmawan, I.S. Surono, Y.K. Lee, Adhesion properties of indigenous dadih lactic acid bacteria on human intestinal mucosal surface, *Asian-Australas. J. Anim. Sci.* 19 (5) (2006) 751–755.
- [33] T. Shi, K. Nishiyama, K. Nakamata, et al., Isolation of potential probiotic *Lactobacillus rhamnosus* strains from traditional fermented mare milk produced in Sumbawa Island of Indonesia, *Biosci. Biotechnol. Biochem.* 76 (10) (2012) 1897–1903.
- [34] I.N. Sujaya, Y. Ramona, N.P. Widarini, et al., Characterization of lactic acid bacteria isolated from Sumbawa mare milk, *J. Vet.* 9 (2) (2008) 52–59.
- [35] N.S. Antara, I.N. Dibia, W.R. Aryanta, Characterization of lactic acid bacteria isolated from horse milk of Bima, *Agritech* 29 (1) (2009) 1–9.
- [36] K.J. Heller, Probiotic bacteria in fermented foods: product characteristics and starter organisms, *Am. J. Clin. Nutr.* 73 (Suppl.) (2001) 374S–379S.
- [37] FAO/WHO [Food and Agriculture Organization/World Health Organization], Guidelines for the Evaluation of Probiotics in Foods, Report of a Joint FAO/WHO Working Group, London, Ontario, Canada, 2002.
- [38] S. Salminen, S. Gorbach, Y. Lee, et al., Human studies on probiotics: what is it really proven today? in: S. Salminen, A. von Wright, A. Ouwehand (Eds.), *Lactic Acid Bacteria: Microbiology and Functional Aspects*, 3rd ed., Marcel Dekker Inc., New York, 2004, pp. 515–530 (revised and expanded).
- [39] A.C. Ouwehand, E.M. Tuomola, S.T. Ikko, et al., Assessment of adhesion properties of novel probiotic strains to human intestinal mucus *Int. J. Food Microbiol.* 64 (2001) 119–126.
- [40] M.T. Liong, P. Shah, Optimization of cholesterol removal by probiotics in the presence of prebiotics by using a response surface method, *Appl. Environ. Microbiol.* 71 (4) (2005) 1745–1753.
- [41] M.P. Díaz-Ropero, R. Martín, S. Sierra, et al., Two *Lactobacillus* strains, isolated from breast milk, differently modulate the immune response, *J. Appl. Microbiol.* 102 (2) (2007) 337–343.
- [42] Y. Kotani, S. Shinkai, H. Okamatsu, et al., Oral intake of *Lactobacillus pentosus* strain b240 accelerates salivary immunoglobulin A secretion in the elderly: a randomized, placebo-controlled, double blind trial, *Immun. Ageing* 7 (11) (2010) 1–11.
- [43] S. Salminen, S. Nybom, J. Meriluoto, et al., Interaction of probiotics and pathogens—benefits to human health? *Curr. Opin. Biotechnol.* 21 (2010) 157–167.
- [44] T. Djaafar, E.S. Rahayu, D. Wibowo, et al., Antimicrobial substance produced by *Lactobacillus* TGR-2 isolated from growol, *Indones. Food Nutr. Prog.* 3 (2) (1996) 29–34.
- [45] N. Kusumawati, B.S.L. Jenie, S. Setyahadi, et al., Selection of indigenous Lactic Acid Bacteria as probiotic strain to reduce cholesterol, *J. Mikrobiol. Indones.* 8 (2) (2003) 39–43.
- [46] L.G. Ooi, M.T. Liong, Cholesterol-lowering effects of probiotics and prebiotics: a review of *in vivo* and *in vitro* findings, *Int. J. Mol. Sci.* 11 (2010) 2499–2522.
- [47] S.K. Nybom, M.C. Collado, I.S. Surono, et al., Effect of glucose in removal of microcystin-LR by viable commercial probiotic strains and strains isolated from dadih fermented milk, *J. Agric. Food Chem.* 56 (2008) 3714–3720.
- [48] Sumarna, Hydrolysis of bioactive isoflavone in soymilk fermented with  $\beta$ -glucosidase producing lactic acid bacteria from local fermented foods of Indonesian, *Malays. J. Microbiol.* 6 (1) (2010) 30–40.
- [49] N. Suhartatik, M.N. Cahyanto, S. Rahardjo, et al., Isolation and identification of lactic acid bacteria producing  $\beta$  glucosidase from Indonesian fermented foods, *Int. Food Res. J.* 21 (3) (2014) 937–942.
- [50] H. Michlmayr, W. Knefel,  $\beta$ -Glucosidase activities of lactic acid bacteria: mechanisms, impact on fermented food and human health, *FEMS Microbiol. Lett.* 352 (2014) 1–10.
- [51] I. Susanti, R.W. Kusumaningtyas, F. Illaningtyas, Probiotic characteristic of lactic acid bacteria as candidate for functional food, *J. Teknol. Ind. Pangan* 18 (2) (2007) 89–95.
- [52] E.S. Rahayu, A. Yogeswara, T. Utami, et al., Indigenous probiotic strains of Indonesia and their application for fermented food, in: *Proceedings of the 12th ASEAN Food Conference*, 2011, pp. 400–404.
- [53] Desniar, I. Rusmana, A. Suwanto, et al., Characterization of lactic acid bacteria isolated from an Indonesian fermented fish (bekasam) and their antimicrobial activity against pathogenic bacteria, *Emir. J. Food Agric.* 25 (6) (2015) 489–494.
- [54] C.Y. Ko, H.T.V. Lin, G.J. Tsai, Gamma-aminobutyric acid production in black soybean milk by *Lactobacillus brevis* FPA 3709 and the antidepressant effect of the fermented product on a forced swimming rat model, *Process Biochem.* 48 (2013) 559–568.
- [55] R.D. Cagno, F. Mazzacane, C.D. Rizzello, et al., Synthesis of isoflavone aglycones and equol in soy milks fermented by food-related lactic acid bacteria and their effect on human intestinal caco-2 cells, *J. Agric. Food Chem.* 58 (2010) 10338–10346.
- [56] L. Edwards-Ingram, P. Gitsham, N. Burton, et al., Genotypic and physiological characterization of *Saccharomyces boulardii*, the probiotic strain of *Saccharomyces cerevisiae*, *Appl. Environ. Microbiol.* 73 (8) (2007) 2458–2467.
- [57] Z. Kurugöl, G. Koturoğlu, Effects of *Saccharomyces boulardii* in children with acute diarrhoea, *Acta Paediatr.* 94 (1) (2005) 44–47.
- [58] M. Guslandi, P. Giollo, P.A. Testoni, A pilot trial of *Saccharomyces boulardii* in ulcerative colitis, *Eur. J. Gastroenterol. Hepatol.* 15 (6) (2003) 697–698.
- [59] J.R. Endres, A. Clewell, K.A. Jade, et al., Safety assessment of a proprietary preparation of a novel probiotic, *Bacillus coagulans*, as a food ingredient, *Food Chem. Toxicol.* 47 (2009) 1231–1238.
- [60] D.R. Mandel, K. Eichas, J. Holmes, *Bacillus coagulans*: a viable adjunct therapy for relieving symptoms of rheumatoid arthritis according to a randomized, controlled trial, *BMC Complement Altern. Med.* 10 (2010) 1.
- [61] H.A. Hong, L.H. Duc, S.M. Cutting, The use of bacterial spore formers as probiotics, *FEMS Microbiol. Rev.* 29 (2005) 813–835.
- [62] L. Nuraida, W. Krusong, Starter cultures, in: J.D. Owens (Ed.), *Indigenous Fermented Foods of Southeast Asia*, CRC Press, 2014, pp. 109–136.
- [63] M.M. Ardhana, G.H. Fleet, The microbial ecology of tape ketan fermentation, *Int. J. Food Microbiol.* 9 (1989) 157–165.
- [64] T. Barus, A. Kristina, A. Yuliandi, Diversity of amylase-producing *Bacillus* spp. from "Tape" (fermented cassava), *HAYATI J. Biosci.* 20 (2) (2013) 94–98.
- [65] U. Pato, I.S. Surono, Koesnandar, et al., Hypocholesterolemic effect of indigenous dadih lactic acid bacteria by deconjugation of bile salts, *Asian-Australas. J. Anim. Sci.* 17 (12) (2004) 1741–1745.
- [66] Yuliawati, Y.D. Jurnal, E. Purwati, et al., The effect of *Pediococcus pentosaceus* on stool frequency, TNF- $\alpha$  level, gut microflora balance in

- diarrhea-induced mice, Indones. J. Gastroenterol. Hepatol. Dig. Endosc. 13 (2012) 97–102.
- [67] Aslinar, Y.D. Jurnal, E. Purwati, et al., Probiotic *Weissella paramesenteroides* on enteropathogenic *E. coli*-induced diarrhea, Paediatr. Indones. 54 (1) (2014) 1–8.
- [68] I.M. Mandomando, E.V. Macte, S. Sanz, et al., Etiology of diarrhea in children younger than 5 years of age admitted in a rural hospital of southern Mozambique, Am. J. Trop. Med. Hyg. 76 (3) (2007) 522–527.
- [69] I.S. Surono, F.P. Koestomo, N. Novitasari, et al., Novel probiotic *Enterococcus faecium* IS-27526 supplementation increased total salivary sIgA level and bodyweight of pre-school children: a pilot study, Anaerobe 17 (2011) 496–500.
- [70] I.S. Surono, P.D. Martono, S. Kameo, et al., Effect of probiotic oitL. plantarum IS-10506 and zinc supplementation on humoral immune response and zinc status of Indonesian pre-schoolchildren, J. Trace Elem. Med. Biol. 28 (4) (2014) 465–469.